

Mark Scheme Summer 2008

AEA

AEA Mathematics (9801)

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at www.edexcel.org.uk.

Summer 2008

Publications Code UA020312

All the material in this publication is copyright

© Edexcel Ltd 2008

Contents

1.	AEA Mathematics Mark Scheme	5
----	-----------------------------	---

①. $a=200, d=-\frac{5}{2}$ $u_n = 0 \Rightarrow 200 - \frac{5}{2}(n-1) = 0$

$\Rightarrow n = 81$

[ALT $S_n = \frac{n}{2}(400 - \frac{5}{2}(n-1))$



Max at 80.5

Identify a, d and set $u_n = 0$

S_n and attempt max

M1
A1
M1
A1]

Maximum sum when $n=80$ or 81

$S_{80} = 40 [400 - \frac{5}{2} \times 79]$
 $= 20 [800 - 395]$
 $= \underline{\underline{8100}}$

Use of S_n with $n=80$ or $n=81$

M1 A1
A1 (5)

②. (a) $\frac{dy}{dx} = 2 \Rightarrow 2(x+1)(x+2) = xy$

$\Rightarrow 2(x^2 + 3x + 2) = x(2x+5)$

$y = 2x+5 \Rightarrow \underline{\underline{x = -4, y = -3}}$ [or P is (-4, -3)]

(b) $\int \frac{1}{y} dy = \int \frac{x}{(x+1)(x+2)} dx$
 $= \int (\frac{2}{x+2} - \frac{1}{x+1}) dx$

$\Rightarrow \ln|y| = 2\ln|x+2| - \ln|x+1| + C$

$\ln y = \ln \left[\frac{A(x+2)^2}{(x+1)} \right] + C$ or $\ln \left[\frac{(x+2)^2}{(x+1)} \right] + C$

$y = \frac{A(x+2)^2}{(x+1)}$

Using P(-4, -3) $\Rightarrow -3 = \frac{A(-2)^2}{(-3)}$ (✓ their P)

$\underline{\underline{y = \frac{9(x+2)^2}{4(x+1)}}}$

Sub $\frac{dy}{dx} = 2$

sub y for $2x+5$ and attempt to solve

Separation attempt

Attempt partial fractions

Some correct ln integral of x function

Use of log rules $\rightarrow \ln[g(x)]$ (condone A=1 or C=0)

Getting out of loss (must have 'A' or equiv)

Use P to form eqn in A or C


M1
M1
A1
A1 (4)

M1
M1
M1 A1
M1
M1
M1
A1 (8)

Max S1 only for 11 or 12/12

(12)

S1 or S2 For a fully correct (or nearly so) and neat or succinct solution to Qn 2 - Qn 7. Count best 3 questions.
I1 For a good attempt at the whole paper (all questions).

③ (a) $\tan 2\theta = \frac{2\tan\theta}{1-\tan^2\theta}$; $t = \tan 15$ 

$\tan 30 = \frac{1}{\sqrt{3}} \Rightarrow \frac{1}{\sqrt{3}} = \frac{2t}{1-t^2}$

$t^2 + 2\sqrt{3}t - 1 = 0 \Rightarrow t = \frac{-2\sqrt{3} \pm \sqrt{12+4}}{2}$

$t = \tan 15 = \underline{2-\sqrt{3}}$ (*)

Use of known t_2 ... BI
 Use ~~tan~~ → MI
 equation in t
 Attempt to solve → t: MI
 [S for considerations ±] AI csp (4)

(b) $\left(\frac{\sin\theta}{2} + \frac{\sqrt{3}\cos\theta}{2}\right)\left(\frac{\sin\theta}{2} - \frac{\sqrt{3}\cos\theta}{2}\right) = \cos^2\theta(1-\sqrt{3})$

$\frac{\sin^2\theta}{4} - \frac{3}{4}\cos^2\theta = \cos^2\theta - \sqrt{3}\cos^2\theta$

$\frac{\sin^2\theta}{4} = \frac{\cos^2\theta}{4}(7-4\sqrt{3})$

$\cos^2\theta = \frac{1}{4(2-\sqrt{3})}$ or $\frac{2+\sqrt{3}}{4}$ or $\tan^2\theta = 7-4\sqrt{3}$ or $\cos 2\theta = \frac{2\sqrt{3}-3}{4-2\sqrt{3}}$

$\tan^2\theta = (2-\sqrt{3})^2$

$\tan\theta = \pm(2-\sqrt{3})$ or $\cos 2\theta = \frac{\sqrt{3}}{2}$

$\tan\theta = 2-\sqrt{3} \Rightarrow \theta = \underline{15, 195}$; $\tan\theta = -(2-\sqrt{3}) \Rightarrow \theta = \underline{165, 345}$

Use of $\sin(A \pm B)$ MI
 Equation in s^2 and c^2 MI
 or c^2 and $\cos 2\theta$
 Attempt $\cos^2\theta$, $\tan^2\theta$ MI
 or $\cos 2\theta$ or $\sin^2\theta$
 AI
 $(2-\sqrt{3})^2 = 7-4\sqrt{3}$ MI
 AI
 AI; AI (8) (12)

④ (a) $\frac{dy}{dx} = -\sin x \ln(\sec x) + \cos x \cdot \tan x$


$y' = 0 \Rightarrow 0 = \sin x (1 - \ln(\sec x))$

[$\sin x = 0 \Rightarrow x = 0 \therefore$ Min at origin]

$\ln \sec x = 1 \Rightarrow \sec x = e ; \therefore \theta \in (\arccos \frac{1}{e}, \frac{1}{e})$

(2) [Rectangle - S]

Use of product rule MI AI
 Take out $\sin x$ factor MI
 [Smokes]
 AI; AI (5)
 For strategies MI
 Attempt parts MI AI

(b) 

$I = \int \cos x \ln(\sec x) dx = \sin x \ln \sec x - \int \sin x \tan x dx$

$I = \sin x \ln \sec x - \int \frac{\sin^2 x}{\cos x} dx = \sin x \ln \sec x - \int (\sec x - \cos x) dx$

$I = \sin x \ln \sec x - \ln|\sec x + \tan x| + \frac{\sin x}{\cos x}$

$S = [I]_0^{\arccos \frac{1}{e}}$

$S = \frac{\sqrt{e^2-1}}{e} - \ln[e + \sqrt{e^2-1}] + \frac{\sqrt{e^2-1}}{e}$

$\text{Area} = 2\left[\frac{1}{e} \arccos \frac{1}{e} - S\right] = \underline{\underline{\frac{2}{e} \arccos \frac{1}{e} + 2\ln(e + \sqrt{e^2-1}) - \frac{4\sqrt{e^2-1}}{e}}}$ (*)

Put $\sin x \tan x$ into integrable form MI
 correct integration AI AI
 Attempt correct limits and $\sin x$ and $\cos x$ in terms of $e, \sqrt{e^2-1}$ etc. MI
 Must have complete integral
 AI csp. (8)
 (13)

5 (i) $(\log_3 p)^2 = \log_3(p^2) \Rightarrow (\log_3 p)^2 = 2 \log_3 p$
 $\Rightarrow \log_3 p (\log_3 p - 2) = 0 \Rightarrow \log_3 p = 0 \therefore p = 1$
 or $\log_3 p = 2 \therefore p = 9$
 $\log_3(p+q) = \log_3 p + \log_3 q \Rightarrow \log_3(p+q) = \log_3(pq)$
 $\Rightarrow p+q = pq$ or $q+q = 9q$
 $\therefore q = \frac{p}{p-1} \quad (p \neq 1)$
 $p = 9 \Rightarrow q = \frac{9}{8}$



Use $n \log x$ rule
 Use of $\log x + \log y$ rule

M1
 A1
 A1
 M1
 A1

(ii) $\log_3 \left[\frac{3x^3 - 23x^2 + 40x}{(3x-8)^2} \right] = 1$
 $\frac{3x^3 - 23x^2 + 40x}{(3x-8)^2} = 3$
 $\frac{x(3x-8)(x-5)}{(3x-8)^2} = 3$
 $x^2 - 5x = 9x - 24 \Rightarrow x^2 - 14x + 24 = 0$
 $(x-2)(x-12) = 0 \Rightarrow x = 2 \text{ or } 12$
 $(x = 9/8 \text{ listed here loses final A1})$
 $\log_3 9 = 2 \text{ o.e.}$

Making q the subject
 [S for $p \neq 1$]
 Seen anywhere
 Use of log rules to form a single log out of logs
 For reducing cubic equation to quadratic
 [$x \neq 8/3$ for S mob]
 3TQ (accused 3TQ)
 (ignore $x=2$ and $9/8$)
 [Smobs for connection]

M1
 A1 (7)
 B1
 M1
 M1
 M1
 A1
 M1 ; A1 (7)
 (14)

6 (a) $yx + 2y = ax + b \Rightarrow x = \frac{b-2y}{y-a} \therefore f^{-1}(x) = \frac{b-2x}{x-a}$
 (b) $f(f(x)) = x \Rightarrow f^{-1}(b) = f(x) ; \therefore a = -2$
 (c) (i) 
 (ii) 
 (d) Normal at P' on $y = f(x-2) + 2$ is: $y = 4(x-2) - 39 + 2$
 $y = 4x - 45$
 curve is symmetric about $y = \frac{1}{2}x$, normal at Q' will be $y = 4x + 45$
 [symmetry is $x \rightarrow -x$ and $y \rightarrow -y$]
 Reversing process
 Normal at Q on $y = f(x)$ is: $y = 4(x+2) + 45 - 2$
 $\therefore y = 4x + 51 \text{ or } k = 51$


Make x the subject
 $f^{-1} = f$
 shape
 $x = -2; y = -2$
 $(\frac{1}{2}, 0), (0, \frac{1}{2})$
 $\rightarrow +2$
 $\uparrow +2$
 both branches
 Use transformations on normal
 Use symmetry on Q'
 Use $f(x+2) - 2$

M1, A1 (2)
 M1; A1 (2)
 B1 (no overlap)
 B1
 B1 (3)
 M1
 M1
 A1 (3)
 M1
 A1
 M1
 A1

[NB: P' is (12, 3)
 P is (10, 1); $b = 32$; Q = (-14, -5)]
 $y - -5 = 4(x - -14)$
 $\rightarrow k = 51$

M1
 A1

M1
 A1 (5)
 (15)

(7) (a)  $\vec{BA} = \begin{pmatrix} -4 \\ 1 \\ -8 \end{pmatrix}$ $\vec{BC} = \begin{pmatrix} 4 \\ 2 \\ -4 \end{pmatrix}$

$\vec{BA} \cdot \vec{BC} = -16 + 2 + 32 = 18$

$|\vec{BA}| = \sqrt{4^2 + 1^2 + 8^2} = 9$, $|\vec{BC}| = \sqrt{4^2 + 2^2 + 4^2} = 6$

$\cos \theta = \frac{18}{9 \times 6} = \frac{1}{3}$

Attempt \vec{BA} and \vec{BC}

M1


Attempt $\vec{BA} \cdot \vec{BC}$

M1

Attempt $|\vec{BA}|$ or $|\vec{BC}|$

M1

AI (4)



Using rhombus idea, $\vec{BX} = \vec{BC} + \frac{2}{3}\vec{BA}$ o.e.

$= \lambda \begin{pmatrix} 4 \\ 8 \\ -28 \end{pmatrix}$ or $\mu \begin{pmatrix} 1 \\ 2 \\ -7 \end{pmatrix}$

eg $3\vec{BC} + 2\vec{BA}$
Any correct ratio

M1, AI


AI

AI c/o.

(4)

Through B $\therefore \underline{\underline{r = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} + t \begin{pmatrix} 1 \\ 2 \\ -7 \end{pmatrix} (*)}}$

(c) $\vec{AC} = \begin{pmatrix} 8 \\ 1 \\ 4 \end{pmatrix}$ $|\vec{AC}| = \sqrt{8^2 + 1^2 + 4^2} = 9 = |\vec{BA}|$



Attempt \vec{AC} and $|\vec{AC}|$
(must say = $|\vec{BA}|$ for AI)

M1 AI c/o (2)

(d) $\therefore ABC$ is isos L_1 has direction $\frac{1}{2}(\vec{AB} + \vec{AC}) = \begin{pmatrix} 6 \\ 0 \\ 6 \end{pmatrix}$

$\therefore L_1$ has equation $(r =) \begin{pmatrix} -3 \\ 1 \\ -9 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$

Centre of S is intersection of L_1 and L

Solving: $\left. \begin{matrix} 1 + t = -3 + u \\ 2t = 1 \end{matrix} \right\} \Rightarrow t = \frac{1}{2}, u = \frac{9}{2}$

$\left[-1 - 7t = -9 + u \right]$ Check: LHS = $-\frac{9}{2}$, RHS = $-\frac{9}{2}$

\therefore Centre has position vector $\underline{\underline{\vec{ON} = \begin{pmatrix} 3/2 \\ 1 \\ -9/2 \end{pmatrix}}}$

Find equation of L_1

BI

M1

Strategy

M1


Attempt to solve $t = \frac{1}{2}, u = \frac{9}{2}$

M1

AI

(-100)

A2/1/0 (7)

(e)  let X = mid-point of BC $\therefore BX = 3$ (\therefore isos)

$\vec{BN} = \begin{pmatrix} 1/2 \\ 1 \\ -7/2 \end{pmatrix}$ $\therefore BN = \frac{1}{2}\sqrt{54}$

$BX = 3$

BI

Attempt \vec{BN} or $|\vec{BN}|$

M1

AI

$r^2 = BN^2 - 3^2 \therefore r^2 = \frac{54}{4} - 9 \therefore r = \frac{\sqrt{18}}{2}$ or $\frac{3\sqrt{2}}{2}$

Full method for r

M1 AI (5)

(22)

Final
CB Afford
1/7/08

3(a) $\tan 15 = \tan(60-45) = \frac{\tan 60 - \tan 45}{1 + \tan 60 \tan 45} = \frac{\sqrt{3} - 1}{1 + \sqrt{3}}$
 $= \frac{(\sqrt{3}-1)^2}{3-1}, = \frac{4-2\sqrt{3}}{2} = \underline{\underline{\frac{2-\sqrt{3}}{1}}}$

or $\tan(45-30)$
 NB no need for \pm consideration this way

BI MI
 MI, AI (4)

3(b) Use of $\cos A - \cos B = \dots = 2 \sin(\frac{20+120}{2}) \sin(\frac{20-120}{2})$
 $\rightarrow -\frac{1}{2} [\cos 2\theta - \cos 120] = (1-\sqrt{3}) \cos^2 \theta$
 $\rightarrow \cos 2\theta + \frac{1}{2} = 2(\sqrt{3}-1) \cos^2 \theta$
 May then go for $\cos^2 \theta = \dots$ or $\cos 2\theta = \dots$ as per scheme.


See snippets
 3A

MI
 MI

6(d) For a valid method $\rightarrow P = (10, 1)$ or $P' = (12, 3)$
 For obtaining $Q = (-14, -5)$
 Equation of normal: $y - 5 = 4(x - 14)$
 $y = 4x + 51$ or $k = 51$

Requires $b = 32$ on the way!

MI AI
 MI
 MI
 AI (5)

7(e)  Midpoint of BC, $\vec{OX} = \begin{pmatrix} 3 \\ -3 \end{pmatrix}$
 $\vec{NX} = \pm \begin{pmatrix} 3/2 \\ 0 \\ 3/2 \end{pmatrix}$
 $r = |\vec{NX}| = \sqrt{(\frac{3}{2})^2 + (\frac{3}{2})^2} = \underline{\underline{\frac{3}{2}\sqrt{2}}}$ o.e.

Must be in (e)

BI
 MI AI
 MI AI (5)

7(d) $\vec{ON} = \begin{pmatrix} 1+t \\ 2t \\ -1-7t \end{pmatrix}$
 $\vec{AN} = \begin{pmatrix} 4+t \\ -1+2t \\ 8-7t \end{pmatrix}$
 $\vec{AN} \cdot \vec{BC} = 0 \Rightarrow 16+4t - 2+4t - 32+28t = 0$
 $36t = 18$
 $t = \frac{1}{2}$

General $\vec{ON} =$

Attempt \vec{AN} in t

$\vec{AN} \cdot \vec{BC} = 0$

Solve for t

Then as before.

BI
 MI
 MI
 MI

2(b) $\ln|y| = \frac{1}{2} \int \frac{2x+3}{x^2+3x+2} dx - \frac{3}{2} \int \frac{1}{(x+\frac{1}{2})^2 + (\frac{1}{2})^2} dx$
 $= \frac{1}{2} \ln|x^2+3x+2| - 3 \ln \left| \frac{x+\frac{3}{2}-\frac{1}{2}}{x+\frac{3}{2}+\frac{1}{2}} \right| + c$
 $= \dots$

sep
 Equiv to PIF

Correct ln integration

etc.

MI
 MI
 MI
 AI

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481

Email publications@linneydirect.com

Order Code UA020312 Summer 2008

For more information on Edexcel qualifications, please visit www.edexcel.org.uk/qualifications

Edexcel Limited. Registered in England and Wales no.4496750
Registered Office: One90 High Holborn, London, WC1V 7BH