

Mark Scheme (Results)

Summer 2014

Pearson Edexcel GCE in Statistics S1R  
(6683/01R)

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at [www.edexcel.com](http://www.edexcel.com).

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

[www.edexcel.com/contactus](http://www.edexcel.com/contactus)

## **Pearson: helping people progress, everywhere**

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2014

Publications Code UA040120

All the material in this publication is copyright

© Pearson Education Ltd 2014

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

## EDEXCEL GCE MATHEMATICS

### General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.
3. Abbreviations


These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
  - ft – follow through
  - the symbol  $\surd$  will be used for correct ft
  - cao – correct answer only
  - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
  - isw – ignore subsequent working
  - awrt – answers which round to
  - SC: special case
  - oe – or equivalent (and appropriate)
  - dep – dependent
  - indep – independent
  - dp decimal places
  - sf significant figures
  - \* The answer is printed on the paper
  - The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.
  5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
  6. If a candidate makes more than one attempt at any question:
    - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
    - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
  7. Ignore wrong working or incorrect statements following a correct answer.

Question Number	Scheme	Marks
<p><b>1. (a)</b></p> $0.4 + p + 0.05 + 0.15 + p = 1$ $2p = 0.4$ $p = 0.2$ <p><b>(b)</b></p> $E(X) = 0.4 \times -4 + 0.2 \times -2 + 0.05 \times 1 + 0.15 \times 3 + 0.2 \times 5 = -0.5$ <p><b>(c)</b></p> $[F(0) = P(X = -2) + P(X = -4) = 0.2 + 0.4] = 0.6$ <p><b>(d)</b></p> $P(3X + 2 > 5) = P(X > 1)$ $P(3X + 2 > 5) = P(X = 3) + P(X = 5)$ $P(3X + 2 > 5) = 0.35$ <p><b>(e)</b></p> $\text{Var}(aX + 3) = a^2 \text{Var}(X)$ $53.4 = a^2 13.35$ $a = \pm 2$	<p><u>or</u> verify <math>0.4 + 0.2 + 0.05 + 0.15 + 0.2 = 1</math></p> <p>conclusion <math>p = 0.2</math> must be stated</p>	<p>M1</p> <p>A1cso (2)</p> <p>M1 A1 (2)</p> <p>B1 (1)</p> <p>M1 A1 (2)</p> <p>M1 A1 (2)</p> <p>[ <b>Total 9</b> ]</p>
<b>Notes</b>		
	<p><b>(a)</b> M1 for equating sum of all probabilities to 1 The minimum working required is: <math>0.6 + 2p = 1</math> but <math>2p = 1 - 0.6</math> or <math>2p = 0.4</math> is M0 BUT allow <math>1 - 0.4 - 0.05 - 0.15 = 0.4</math> followed by <math>2p = 0.4</math> <u>or</u> <math>1 - 0.4 - 0.05 - 0.15 = 2p</math> Since <u>all</u> of the probabilities are seen.</p> <p>A1cso for a correct solution with no incorrect working seen (For verify method, they must conclude that <math>p = 0.2</math>)</p> <p><b>(b)</b> M1 for a correct expression with at least 3 correct terms May be: <math>-1.6 - 0.4 + 0.05 + 0.45 + 1</math> A1 for <math>-0.5</math></p> <p><b>(c)</b> B1 for 0.6</p> <p><b>(d)</b> M1 for identifying <math>X = 3</math> and <math>X = 5</math> only (<math>X &gt; 1</math> is not sufficient) A1 for 0.35</p> <p><b>(e)</b> M1 for <math>\text{Var}(aX + 3) = a^2 \text{Var}(X)</math> but this may be implied by seeing <math>a = 2</math> <u>or</u> <math>a = -2</math> A1 for <u>both</u> correct values <math>+2</math> <u>and</u> <math>-2</math></p>	


Question Number	Scheme	Marks
<p>2. (a)</p> <p>(b)</p> <p>(c)</p>	<p>(Discrete) Uniform</p> <p>(i) <math>P(X = 10) = \frac{1}{10}</math></p> <p>(ii) <math>P(X &lt; 10) = \frac{9}{10}</math></p> <p>(i) <math>P(Y = 10) = 0</math></p> <p>(ii) <math>P(Y &lt; 10) = \frac{1}{2}</math></p>	<p>B1</p> <p>(1)</p> <p>B1</p> <p>B1</p> <p>(2)</p> <p>B1</p> <p>B1</p> <p>(2)</p> <p>[ Total 5]</p>
	<b>Notes</b>	
(a)	<p>B1 for seeing the word <b>uniform</b></p> <p>Condone “continuous” uniform</p>	


Question Number	Scheme	Marks
<p><b>3. (a)</b></p> <p><math>\sum x = 88</math></p> <p><math>S_{pp} = 11422 - \frac{299.2^2}{8} = [231.92] \quad (*)</math></p> <p><math>S_{xx} = 1160 - \frac{88^2}{8} = 192</math></p> <p><math>S_{xp} = 3449.5 - \frac{88 \times 299.2}{8} = 158.3 \quad \text{awrt } 158</math></p> <p><b>(b)</b></p> <p><math>r = \left[ \frac{S_{xp}}{\sqrt{S_{xx} S_{pp}}} \right] = \frac{158.3}{\sqrt{192 \times 231.92}}</math></p> <p><math>r = 0.7501726031\dots \quad \text{awrt } 0.750</math></p> <p><b>(c)</b></p> <p><math>b = \left[ \frac{S_{xp}}{S_{xx}} \right] = \frac{158.3}{192} = 0.824(479166\dots) \quad (*)</math></p> <p><math>a = \bar{p} - b\bar{x} = \frac{299.2}{8} - 0.824\dots \times \frac{88}{8} = 28.330729\dots \quad \text{awrt } 28.3</math></p> <p><b>(d)</b> <math>p = 28.3\dots + 0.824\dots \times 10 = 36.57552\dots \quad \text{awrt } \text{£}3700</math></p> <p><b>(e)</b> Goes up £82.40</p> <p><b>(f)</b> (i) <math>r = 0.750</math> (ii) <math>b = 0.412</math></p>		<p>B1</p> <p>B1cso</p> <p>M1 A1</p> <p>A1</p> <p><b>(5)</b></p> <p>M1</p> <p>A1</p> <p><b>(2)</b></p> <p>M1 A1cso</p> <p>M1 A1</p> <p><b>(4)</b></p> <p>M1 A1 <b>(2)</b></p> <p>B1</p> <p><b>(1)</b></p> <p>B1ft</p> <p>B1 <b>(2)</b></p> <p><b>[Total 16]</b></p>
<b>Notes</b>		
	<p><b>(a)</b> 1<sup>st</sup> B1 for <math>\sum x = 88</math> seen. May be in a correct formula or implied by 192 or 158.3  2<sup>nd</sup> B1cso for a correct expression for <math>S_{pp}</math>  M1 for a correct expression for <math>S_{xx}</math> or <math>S_{xp}</math> (ft their <math>\Sigma x</math>). If we don't see an explicit <math>\Sigma x = k</math> but consistent use of <math>k</math> instead of 88 in <math>S_{xp}</math> <u>and</u> <math>S_{xx}</math> then award M1  1<sup>st</sup> A1 for <math>S_{xx} = 192</math>      2<sup>nd</sup> A1 for <math>S_{xp} = \text{awrt } 158</math></p> <p><b>(b)</b> M1 for correct expression for <math>r</math> ft their 192 and 158.3 May be implied by <math>r = 0.75</math>  A1 for awrt 0.750 Allow A1 for <math>r = 0.75</math> if a correct expr' is seen (since 3<sup>rd</sup> sf is 0)</p> <p><b>(c)</b> 1<sup>st</sup> M1 for a correct expression for <math>b</math> using their values NB. use of 158 gives 0.8229  1<sup>st</sup> A1 cso for <math>b = \text{awrt } 0.824</math></p> <p><b>SC</b> If there is no expression but 0.8244...or better is seen award 1 mark as M0A1  2<sup>nd</sup> M1 for a correct expression for <math>a</math> ft their <math>\Sigma x</math></p> <p><b>(d)</b> 2<sup>nd</sup> A1 for <math>a = \text{awrt } 28.3</math>  M1 for substituting <math>x = 10</math> into their equation</p> <p><b>(e)</b> A1 for awrt £3700 (£ 36.58 or £36.58 (hundreds) is A0)</p> <p><b>(f)(i)</b> B1 for goes up £82.40 (for each additional employee) (£0.824 hundreds is B0)</p> <p><b>(ii)</b> B1ft for <math>r =</math> their answer to (b) . Allow recalculation. Condone <math> r  &gt; 1</math>  B1 for 0.412 only</p>	

Question Number	Scheme	Marks
<p>4. (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p><math>P(A \cap B) = P(A B) \times P(B)</math></p> <p><math>P(A \cap B) = \frac{2}{5} \times \frac{1}{2} = \frac{1}{5}</math></p> <div style="border: 1px solid black; padding: 10px; display: inline-block; margin: 10px;">  </div> <p>2 intersecting circles and 'P(A ∩ B)'</p> <p><math>\frac{3}{20}</math> and <math>\frac{3}{10}</math></p> <p>Box and <math>\frac{7}{20}</math></p> <p><math>\left[ P(A) = \frac{3}{20} + \frac{1}{5} \right] = \frac{7}{20}</math> or 0.35</p> <p><math>P(B A) = \frac{P(A \cap B)}{P(A)} = \frac{\frac{1}{5}}{\frac{7}{20}} = \frac{4}{7}</math></p> <p>0.3</p>	<p>M1 A1</p> <p>(2)</p> <p>B1ft</p> <p>B1</p> <p>B1</p> <p>(3)</p> <p>B1ft</p> <p>(1)</p> <p>M1</p> <p>A1 cao</p> <p>(2)</p> <p>B1ft</p> <p>(1)</p> <p>[Total 9]</p>
<b>Notes</b>		
	<p>(a) M1 for <math>\frac{2}{5} \times \frac{1}{2}</math> <u>or</u> a correct probability product expression and one correct prob. Ans only 2/2</p> <p>(b) 1<sup>st</sup> B1 for 2 intersecting circles labelled A and B and ft their prob. for intersection Condone missing labels for 2<sup>nd</sup> and 3<sup>rd</sup> B marks</p> <p>(c) B1ft for 0.35 (o.e.) if no Venn diagram <u>or</u> correct follow through from their diagram <u>or</u> allow 0.35 (or correct ft) from correct working e.g. <math>0.65 - 0.5 + (a)</math> B0 for 0.35 if their diagram does not give 0.35 unless it comes from correct work Don't insist on <math>P(A) = \dots</math> but do not award for <math>P(A' \cap B') = \frac{7}{20}</math></p> <p>(d) M1 for <math>\frac{\text{their (a)}}{\text{their (c)}}</math> <u>or</u> a correct ratio of probabilities from their diagram NB incorrect use of <math>P(A' \cap B') = \frac{7}{20}</math> scores M0 and num <math>\geq</math> denom scores M0 A1 for <math>\frac{4}{7}</math> only</p> <p>(e) B1ft for 0.3 <u>or</u> correct ft from their Venn diagram <u>or</u> ft from <math>\frac{13}{20} - \text{their (c)}</math></p>	



Question Number	Scheme	Marks														
<p>5. (a)</p> <p>(b)</p> <table border="1" data-bbox="349 294 678 541"> <thead> <tr> <th>Time</th> <th>Frequency density</th> </tr> </thead> <tbody> <tr><td>2-4</td><td>5</td></tr> <tr><td>5-6</td><td>4.5</td></tr> <tr><td>7</td><td>6</td></tr> <tr><td>8</td><td>24</td></tr> <tr><td>9-10</td><td>7</td></tr> <tr><td>11-15</td><td>2.4</td></tr> </tbody> </table> <p>(c)</p> <p>(d)</p> <p>(e)</p>	Time	Frequency density	2-4	5	5-6	4.5	7	6	8	24	9-10	7	11-15	2.4	<p>4.5</p> <p>f.d = 24 is represented as 6cm, so f.d. = 7 is represented as <b>1.75(cm)</b></p> <p><math>\frac{1}{3} \times 15 + 9 + \frac{1}{2} \times 6 = 17</math></p> <p>Median = <math>7.5 + \frac{40-30}{24} \times 1 = 7.91666\dots</math> <b>awrt</b> 7.92 or 7.93(75)</p> <p><math>Q_1 = 4.5 + \frac{20-15}{9} \times 2 = 5.611111\dots</math> <b>awrt</b> 5.61 or 5.66(666...)</p> <p><math>Q_3 = 8.5 + \frac{60-54}{14} \times 2 = 9.357142\dots</math> <b>awrt</b> 9.36 or 9.46(4285...)</p> <p><math>Q_3 - Q_2 (= 1.4 \text{ or } 1.5) &lt; Q_2 - Q_1 (= 2.3)</math> <u>or</u> (Mean) &lt; Median &lt; Mode Therefore negative skew</p>	<p>B1 (1)</p> <p>M1 A1</p> <p>A1 (3)</p> <p>M1, A1 (2)</p> <p>M1 A1 A1 A1 (4)</p> <p>B1ft dB1cao (2)</p> <p><b>[Total 12]</b></p>
Time	Frequency density															
2-4	5															
5-6	4.5															
7	6															
8	24															
9-10	7															
11-15	2.4															
<b>Notes</b>																
	<p>(a) B1 for 4.5 (o.e.) only. NB 1.5~4.5 is B0</p> <p>(b) M1 for evidence of f/w (at least 3 f.d. found). May be implied by a correct answer. A1 for identifying 9-10 as 2<sup>nd</sup> highest bar from correct working e.g. <math>24x = 6 \times 7</math> A1 for 1.75(cm). Correct answer only 3/3</p> <p>(c) M1 for a correct expression. May interpolate e.g. <math>[24 + \frac{1}{2} \times 6 - \frac{2}{3} \times 15]</math> or <math>(27 - 10)</math> A1 for 17</p> <p>(d) M1 for one correct fraction in an expression for <math>Q_1</math>, <math>Q_2</math> or <math>Q_3</math> 1<sup>st</sup> A1 for <math>Q_2</math> <b>awrt</b> 7.92 (or 7.94 if <math>(n+1)</math> used – look for 40.5 instead of 40) 2<sup>nd</sup> A1 for <math>Q_1</math> <b>awrt</b> 5.61 (or 5.67 if <math>(n+1)</math> used – look for 20.25 instead of 20) 3<sup>rd</sup> A1 for <math>Q_3</math> <b>awrt</b> 9.36 (or 9.46 if <math>(n+1)</math> used – look for 60.75 instead of 60)  NB watch out for working down e.g. <math>8.5 - \frac{14}{24} \times 1</math> for <math>Q_2</math></p> <p>(e) 1<sup>st</sup> B1ft for a correct comparison of their quartiles e.g. <math>Q_2</math> closer to <math>Q_3</math> or using at least two of Mean &lt; Median &lt; Mode (must state mean or mode if using this method). N.B. Mean = 7.71875, mode = 8 2<sup>nd</sup> B1cao dependent on 1<sup>st</sup> B1 being awarded for <b>negative skew</b> only (no ft)</p>															

Question Number	Scheme	Marks
<p><b>6. (a)</b></p> <p><b>(b)</b> 15%</p> <p><b>(c)(i)</b></p> <p><b>(ii)</b></p> <p><b>(d)</b></p>	<div style="text-align: center;">  </div> <p>24 and 28 (above the mean)</p> <p>For 0.80 and 0.05 (clearly indicated)</p> <p><math>\frac{(28 - \mu)}{\sigma} = 1.64(49)</math> or <math>\frac{(24 - \mu)}{\sigma} = 0.84(16)</math></p> <p>0.8416 and 1.6449 seen</p> <p><math>\mu = 28 - 1.64(49)\sigma</math> , <math>\mu = 24 - 0.84(16)\sigma</math></p> <p>eliminating <math>\mu</math> or <math>\sigma</math></p> <p><math>\sigma = 4.9794597\dots</math> <b>awrt 4.98</b></p> <p><math>\mu = 19.809286\dots</math> <b>awrt 19.8</b></p> <p><math>z = \frac{(12 - '19.8...')}{'4.97...'}</math></p> <p><math>P(Z &lt; -1.57) = 1 - P(Z &lt; 1.57)</math></p> <p><math>1 - 0.9418 = 0.0582</math> <b>awrt 0.06</b></p>	<p>B1</p> <p>B1</p> <p>(2)</p> <p>B1</p> <p>(1)</p> <p>M1</p> <p>B1</p> <p>A1,A1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>(7)</p> <p>M1</p> <p>dM1</p> <p>A1</p> <p>(3)</p> <p><b>[Total 13]</b></p>
<b>Notes</b>		
<p><b>(a)</b></p> <p><b>(b)</b></p> <p><b>(c)</b></p> <p><b>SC</b></p> <p><b>(d)</b></p>	<p>1<sup>st</sup> B1 24 and 28 labelled on the horizontal axis above the mean in the correct order. They must clearly indicate where 24 and 28 are on the horizontal axis.</p> <p>2<sup>nd</sup> B1 for clear, correct labelling of probabilities. Must be associated with correct <u>area</u>.</p> <p>B1 for 15% or 0.15 NB 0.15% is B0</p> <p>1<sup>st</sup> M1 for <math>\frac{\pm(28 - \mu)}{\sigma} = z_1</math> or <math>\frac{\pm(24 - \mu)}{\sigma} = z_2</math> where <math> z_1  &gt; 1.5</math> and <math> z_2  &lt; 1</math></p> <p>Condone <math>z_2 = 0.8</math></p> <p>B1 for both values 0.8416 <u>and</u> 1.6449 or better seen. Calc: 0.8416212..., 1.644853..</p> <p>1<sup>st</sup> A1 for <math>\mu = 28 - 1.64(49)\sigma</math> or any correct arrangement (allow 1.64 ~1.65 inclusive)</p> <p>2<sup>nd</sup> A1 for <math>\mu = 24 - 0.84(16)\sigma</math> or any correct arrangement (allow 0.84 or better)</p> <p>2<sup>nd</sup> M1 for an attempt to solve simultaneous equations by eliminating <math>\mu</math> or <math>\sigma</math></p> <p>3<sup>rd</sup> A1 for <b>awrt 4.98</b> (Condone <math>\sigma = 5</math> or awrt 5.0 if B0 scored)</p> <p>4<sup>th</sup> A1 for <b>awrt 19.8</b></p> <p>For use of 0.84 and 1.64 giving <math>\sigma = 5</math> and <math>\mu =</math> awrt 19.8 score M1B0A1A1M1A1A1</p> <p>or 0.84 and 1.65 giving <math>\sigma =</math> awrt 4.94 and <math>\mu =</math> awrt 19.9 score M1B0A1A1M1A1A1</p> <p>1<sup>st</sup> M1 for standardising with 12, their <math>\mu</math> and <math>\sigma</math> provided <math>\sigma &gt; 0</math></p> <p>If <math>\sigma &lt; 0</math> from their equations in (c) allow M1 if they use <math> \sigma </math></p> <p>2<sup>nd</sup> dM1 for <math>1 - P(Z &lt; '1.57')</math> dependent on the 1<sup>st</sup> M1 being scored i.e. leads to prob <math>&lt; 0.5</math></p> <p>A1 for awrt 0.06 from correct working</p>	

Question Number	Scheme	Marks
<p><b>7. (a)</b></p> <p><b>(b)</b></p> <p><b>(c)</b></p> <p><b>(d)</b></p>	 <p><b>(b)</b> <math>P(\text{car or bike but not both}) = 0.78 \times 0.70 + 0.22 \times 0.85 = 0.733</math></p> <p><b>(c)</b> <math>[P(\text{car} \text{bike})] = \frac{P(\text{car} \cap \text{bike})}{P(\text{bike})} = \frac{0.78 \times 0.30}{0.78 \times 0.30 + 0.22 \times 0.85}, = 0.555819....</math> awrt 0.556</p> <p><b>(d)</b> <math>P(\text{bike}) = 0.78 \times 0.30 + 0.22 \times 0.85 = 0.421</math>, <math>P(\text{not bike}) = 1 - 0.421</math> <math>0.421 \times 0.579 + 0.579 \times 0.421</math> <math>= 0.487518</math> awrt 0.488</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>(3)</p> <p>M1 A1</p> <p>(2)</p> <p>M1A1</p> <p>A1</p> <p>(3)</p> <p>M1</p> <p>dM1</p> <p>A1 (3)</p> <p>[Total 11]</p>
<b>Notes</b>		
<p><b>(a)</b></p> <p><b>(b)</b></p> <p><b>(c)</b></p> <p><b>(d)</b></p>	<p>1<sup>st</sup> B1 for a (2+4) tree with 6 branches</p> <p>2<sup>nd</sup> B1 for 0.78 with label</p> <p>3<sup>rd</sup> B1 for 0.30 and 0.85 with label</p> <p>M1 for correct expression of follow through their correct tree branches</p> <p>A1 for 0.733 or exact equivalent e.g. <math>\frac{733}{1000}</math> and allow 73.3%</p> <p>M1 for a correct expression correct ft <u>or</u> correct formula and <math>\frac{1 \text{ product}}{\text{sum of 2 products}}</math></p> <p>With at least 2 products correct or correct ft. Ratio must be smaller than 1</p> <p>1<sup>st</sup> A1 for finding the denominator correctly. Fully correct expression or = 0.421 (oe)</p> <p>2<sup>nd</sup> A1 for awrt 0.556 or exact equivalent e.g. <math>\frac{234}{421}</math> and allow 55.6%</p> <p>M1 for their <math>P(\text{bike}) \times (1 - P(\text{bike}))</math></p> <p>dM1 for <math>\times 2</math></p> <p>A1 for awrt 0.488</p>	

